

**BOUNDED, ADJOINTABLE OPERATORS ON DENSE SUBSPACES
OF SEPARABLE HILBERT SPACES**

A Thesis

Submitted to the Faculty

in partial fulfillment of the requirements for the

degree of

Bachelor of Arts

in

Mathematics

by

Varun Malladi

Advisor

John D. Trout

DARTMOUTH COLLEGE

Hanover, New Hampshire

May 2023

Abstract

Among inner product spaces, separable Hilbert spaces stand out in their amenability. In this paper, we begin considering how the theory of separable Hilbert spaces influences certain incomplete inner product spaces W , namely those that arise as proper dense subspaces of a separable Hilbert space \mathcal{H} . By the Riesz-Fréchet theorem, every bounded operator $T \in B(\mathcal{H})$ possesses an adjoint $T^* \in B(\mathcal{H})$. The existence of an adjoint is not guaranteed for bounded operators on W ; rather than studying the algebra $B(W)$ of bounded operators on W , we instead restrict our focus to the algebra $BA(W)$ of bounded, *adjointable* operators on W . This is a $*$ -subalgebra of $B(\mathcal{H})$, whose operator norm closure $\overline{BA(W)} := C^*(W)$ is a C^* -subalgebra of $B(\mathcal{H})$. The main focus of this paper is to determine whether $C^*(W) = B(\mathcal{H})$. We find this to be true when W is the span of an orthonormal basis for \mathcal{H} , and provide an example where this is false when W is not the span of an orthonormal basis.

Acknowledgments

I would like to thank my advisor, Professor Jody Trout. Through this experience I have become enchanted by operator algebras, captivated by analysis, and have fallen in love with math all over again. I want to thank Professor Dana Williams, for his encouragement and for connecting me with Professor Trout. I would also like to thank Professor John Voight, for providing feedback on my initial ideas.

Lastly, none of this would be possible without the support of my family and friends. Thank you.

Contents

- Abstract ii
- Acknowledgments iii

- 1 Motivation and Summary 1**
 - 1.1 Introduction 1
 - 1.2 Summary of Results 2

- 2 Linear and Algebraic Structure 4**
 - 2.1 Linear Structure 4
 - 2.2 The Algebra $BA(W)$ 6

- 3 Spectral Theory 11**
 - 3.1 Introduction 11
 - 3.2 On Dense Subspaces 15
 - 3.3 Permanence of Approximate Eigenvalues 17
 - 3.4 Spectral Permanence for Normal Operators 18

- 4 Approximation Results 20**
 - 4.1 Approximation of Compact Operators 20
 - 4.1.1 Finite Rank Operators 21
 - 4.1.2 Projection Operators 23

4.1.3	Approximation Result	27
4.2	Operators on the Span of an ON basis	27
4.2.1	Compatibility $BA(W_0)$ and $BA(W)$	27
4.2.2	Density of $BA(W_0) \subset B(\mathcal{H})$	28
4.3	Example: $BA(W) \subset B(\mathcal{H})$ not Dense	32
4.3.1	An Orthogonal Sequence: (ξ_n)	32
4.3.2	The Operator V	34
4.3.3	Two Sequences	38
4.3.4	Failure to Approximate V	39
5	Future Directions	42
	References	44

Chapter 1

Motivation and Summary

Section 1.1

Introduction

On a Hilbert space \mathcal{H} , every bounded operator $T \in B(\mathcal{H})$ possesses an adjoint $T^* \in B(\mathcal{H})$ by the Riesz-Fréchet theorem. This property is of great utility in the theory, and settings in which bounded operators fail to have an adjoint are generally more difficult to analyze. In these situations, we ask how much of the Hilbert space theory can be recovered by restricting our attention to bounded, adjointable operators, i.e. bounded operators whose adjoint exists.

In this paper, we will consider certain subspaces $W \subset \mathcal{H}$. It is precisely because W may fail to be complete that its bounded operators may fail to have an adjoint. However, there are generalizations when, roughly speaking, completeness of an “inner product” space is not enough to ensure the existence of adjoints. For example, a *Hilbert module* is a (complex) vector space which is complete in the norm defined by a certain sesquilinear form taking values in a C^* -algebra (acting like a generalized inner product). It is already commonplace to restrict study from bounded operators to bounded, adjointable operators in this situation. Taking inspiration from this, our goal is to apply this perspective to proper dense subspaces

of Hilbert spaces.

Specifically, we will consider proper, dense subspaces of separable Hilbert spaces. For the rest of this paper, unless otherwise specified, let \mathcal{H} be a separable, infinite-dimensional, complex Hilbert space and $W \subset \mathcal{H}$ be a proper dense subspace. Let $B(\mathcal{H})$ denote the (C^* -algebra of) bounded operators on \mathcal{H} . We say call a bounded operator $T : W \rightarrow W$ *adjointable* on W if there exists a bounded operator $T^* : W \rightarrow W$ such that $\langle Tx, y \rangle = \langle x, T^*y \rangle$ for all $x, y \in W$. The set of all bounded, adjointable operators on W is denoted as $BA(W)$.

Section 1.2

Summary of Results

We begin by showing compatibility between $BA(W)$ and $B(\mathcal{H})$, for example:

- (a) Every operator $T \in BA(W)$ extends uniquely to an operator $\tilde{T} \in B(\mathcal{H})$.
- (b) Adjoints extend to adjoints: for $T \in BA(W)$, $\tilde{T}^* = (\tilde{T})^*$.
- (c) Normal operators extend to normal operators: if $T \in BA(W)$ is normal, then $\tilde{T} \in B(\mathcal{H})$ is normal.

We then consider how the spectra between T and \tilde{T} differ. By the spectrum of $T \in BA(W)$ we mean the operator spectrum $\sigma(T)$, which we define and show to be distinct from the algebraic spectrum $\sigma_{BA(W)}(T)$. For operators in $B(\mathcal{H})$, the operator spectrum and algebraic spectrum coincide. In both cases, we can partition the spectrum into the point, continuous, and residual spectra, i.e. $\sigma(T) = \sigma_p(T) \sqcup \sigma_c(T) \sqcup \sigma_r(T)$. The notion of approximate eigenvalue $\lambda \in \sigma_{ap}(T)$ also makes sense in both cases. We show the following:

- (a) $\sigma(T) \supset \sigma(\tilde{T})$.
- (b) $\sigma_p(T) \subset \sigma_p(\tilde{T})$.

(c) $\sigma_{ap}(T) = \sigma_{ap}(\tilde{T})$.

(d) If T is normal, then $\sigma_{ap}(T) = \sigma(\tilde{T})$.

Next, we observe the non-functoriality of $BA(-)$: given a sequence of subspace inclusions $W_0 \subset W \subset \mathcal{H}$, it is possible that

(a) $BA(W_0) \not\subset BA(W)$.

(b) $BA(W) \not\subset BA(W_0)$.

Nonetheless, the closure $C^*(W)$ for any dense subspace W always contains the compact operators, i.e. $K(\mathcal{H}) \subset C^*(W)$.

Finally, we consider the question of whether $BA(W) \subset B(\mathcal{H})$ is dense. Our main results are as follows:

(a) If W is the span of an orthonormal basis, then $BA(W) \subset B(\mathcal{H})$ is dense.

(b) As a counterexample of where this fails to hold when W is not the span of an orthonormal basis, $BA(\ell^1)$ is not dense in $B(\ell^2)$.

Chapter 2

Linear and Algebraic Structure

Section 2.1

Linear Structure

To begin, we recall the following:

Lemma 2.1.1. A countable orthonormal basis $\{e_n\}_{n \in \mathbb{N}} \subset \mathcal{H}$ is a collection of orthonormal vectors that is *complete*. Completeness in this situation can be characterized in the following equivalent ways:

- for any vector y , if $\langle y, e_n \rangle = 0$ for all $n \in \mathbb{N}$, then $y = 0$
- $\text{span}\{e_n\}_{n \in \mathbb{N}}$ is dense

If \mathcal{H} possesses a countable orthonormal basis, then

$$x = \sum_{n=1}^{\infty} \langle x, e_n \rangle e_n.$$

for any $x \in \mathcal{H}$.

Proof. See, for example, Theorem 4.14 and Theorem 4.15 in [10]. □

Lemma 2.1.2. A Hilbert space is separable (in the topological sense) if and only if it has a countable, orthonormal basis.

Proof. See, for example, Chapter 6.4 Theorem 9' in [5]. □

The following allows us to consider such a basis to be contained in W :

Proposition 2.1.3. There exists a countable orthonormal basis for \mathcal{H} contained in W .

Proof. Since \mathcal{H} is separable (in the topological sense), and $W \subset \mathcal{H}$ is a proper dense subset, it follows that W is separable. By definition, this means that W contains a countable dense subset $\{a_n\} \subset W$. By applying the Gram-Schmidt process to the linearly independent elements in this subset, we obtain an orthonormal sequence $\{e_n\} \subset W$. Now

$$\text{span}\{a_n\} = \text{span}\{e_n\} \subset W,$$

and by taking closures we see that $\text{span}\{e_n\}$ is dense in \mathcal{H} . By Lemma 2.1.1, this shows that $\{e_n\}$ is a countable orthonormal basis. □

We now state basic facts in Hilbert space theory which still hold on dense subspaces.

Proposition 2.1.4. Let $T \in BA(W)$. Then $\text{Im}(T)^\perp = \text{Ker}(T^*)$.

Proof. First we show $\text{Im}(T)^\perp \subset \text{Ker}(T^*)$. Let $y \in \text{Im}(T)^\perp$. Then, for any $x \in W$, we have $\langle T(x), y \rangle = 0$. Then $\langle x, T^*(y) \rangle = \langle T(x), y \rangle = 0$. By the nondegeneracy of the inner product, it follows that $T^*(y) = 0$. Hence $y \in \text{Ker}(T^*)$.

Showing $\text{Ker}(T^*) \subset \text{Im}(T)^\perp$ is analagous. Let $y \in \text{Ker}(T^*)$. Then $\langle x, T^*(y) \rangle = 0$ for all $x \in W$, and hence $\langle T(x), y \rangle = 0$ for all $x \in W$, which implies $y \in \text{Im}(T)^\perp$. □

Proposition 2.1.5. If $T \in BA(W)$ is normal, then T and T^* have the same kernel.

Proof. We will show $\text{Ker}(T) \subset \text{Ker}(T^*)$, and the other direction is completely analogous. Let $x \in \text{Ker}(T)$. Then $T(x) = 0$, and since $T^*T(x) = TT^*(x) = 0$, it follows that $T^*(x) \in \text{Ker}(T)$.

Now let $y \in W$. We claim that $\langle y, T^*(x) \rangle = 0$, and hence $T^*(x) = 0$ implying $x \in \text{Ker}(T^*)$. To prove the claim, first suppose $y \in \text{Ker}(T)$. Then $\langle y, T^*(x) \rangle = \langle T(y), x \rangle = 0$. On the other hand, suppose $y \notin \text{Ker}(T)$. Recall that the kernel of a continuous linear operator on a normed vector space is closed, so $\text{Ker}(T)$ is closed. But then y is orthogonal to $\text{Ker}(T)$, and since $T^*(x) \in \text{Ker}(T)$ by the previous paragraph, it follows that $\langle y, T^*(x) \rangle = 0$. \square

Section 2.2

The Algebra $BA(W)$

Recall that the set $B(\mathcal{H})$ of bounded linear operators $\mathcal{H} \rightarrow \mathcal{H}$ naturally possesses the structure of a C^* -algebra by taking the $*$ -operation to be “taking adjoints” and the topology to be the operator norm topology. Furthermore, every closed subalgebra of $B(\mathcal{H})$ which is also closed under adjoints is also a C^* -algebra (see the remarks preceding Theorem 4.27 in [2]).

Proposition 2.2.1. Any $T \in B(W)$ has a unique extension $\tilde{T} \in B(\mathcal{H})$. Moreover, $\|T\| = \|\tilde{T}\|$.

Proof. We begin by defining this extension. Let $x \in \mathcal{H}$. Since $W \subset \mathcal{H}$ is dense, there exists a sequence $\{x_n\}$ converging (in \mathcal{H}) to x . We claim $\{T(x_n)\}$ is also a convergent sequence. To see this, note that, for all $n, m > N$,

$$\|T(x_n) - T(x_m)\| = \|T(x_n - x_m)\| \leq \|T\| \|x_n - x_m\|,$$

which shows that $\{T(x_n)\}$ is Cauchy, hence convergent by the completeness of Hilbert spaces.

Finally, define

$$\tilde{T}(x) = \lim_{n \rightarrow \infty} T(x_n).$$

We now verify the validity of this definition.

Well-defined: Write $y = \lim_{n \rightarrow \infty} T(x_n)$. We need to show that if $\{x'_n\} \subset W$ is another sequence converging to x , then $\lim_{n \rightarrow \infty} T(x'_n) = y$. Since $x_n - x'_n \rightarrow 0$, we know from 2.2 that $\|T(x_n) - T(x'_n)\| \rightarrow 0$, and so $T(x_n) - T(x'_n) \rightarrow 0$. This implies $\lim_{n \rightarrow \infty} T(x'_n) = \lim_{n \rightarrow \infty} T(x_n) = y$ as desired.

An extension: Given $v \in W$, we wish to show $\tilde{T}(v) = T(v)$. But this is immediate if we just take the sequence $\{x_n\}$ in the definition above to be the constant sequence $\{v_n\}$, where each $v_n = v$.

Linear: Follows from the linearity of limits.

Bounded: To see this, calculate

$$\begin{aligned} \|\tilde{T}(x)\| &= \left\| \lim_{n \rightarrow \infty} T(x_n) \right\| = \lim_{n \rightarrow \infty} \|T(x_n)\| \\ &\leq \lim_{n \rightarrow \infty} \|T\| \|x_n\| = \|T\| \lim_{n \rightarrow \infty} \|x_n\| \\ &= \|T\| \|x\| < \infty, \end{aligned}$$

where we have repeatedly used the continuity of the norm.

Norm preserving: The above calculation shows that $\|\tilde{T}\| \leq \|T\|$. In detail, let

$$C = \{c \geq 0 : \|\tilde{T}(x)\| \leq c\|x\| \ \forall x \in \mathcal{H}\}.$$

Then $\|T\| \in C$, but $\tilde{T} = \inf(C)$. Now, let

$$C' = \{c \geq 0 : \|T(v)\| \leq c\|v\| \ \forall v \in W\}.$$

Since \tilde{T} is an extension of T , $C \subset C'$. Thus $\|T\| \leq \|\tilde{T}\|$, since

$$\|T\| = \inf(C') \leq \inf(C) = \|\tilde{T}\|.$$

Combining both inequalities gives us $\|T\| = \|\tilde{T}\|$. □

Corollary 2.2.2. The C^* -algebra structure on $B(\mathcal{H})$ determines a natural (unital) $*$ -algebra structure on $BA(W)$.

Corollary 2.2.3. The completion $C^*(W) := \overline{BA(W)} \subset B(\mathcal{H})$ is a C^* -algebra.

Proposition 2.2.4 (properties of extensions). For $T \in B(W)$, let T' be its unique extension to \mathcal{H} . Then:

- (a) $(T - \lambda I)' = T' - \lambda I$.
- (b) $(T^*)' = (T')^*$.
- (c) If $T^{-1} \in B(W)$, then $(T^{-1})' = (T')^{-1}$.

Proof.

- (a) Let $x \in \mathcal{H}$ and $\{w_n\} \subset W$ a sequence converging to x . Then

$$(T - \lambda I)'(x) = \lim_{n \rightarrow \infty} (T - \lambda I)(w_n) = \lim_{n \rightarrow \infty} (T(w_n) - \lambda w_n) = T'(x) - \lambda x = (T' - \lambda I)(x).$$

- (b) It suffices to show that for any $x, y \in \mathcal{H}$, we have

$$\langle T'(x), y \rangle = \langle x, (T^*)'(y) \rangle.$$

Let $\{x_n\}, \{y_n\} \subset W$ be sequences converging to x and y , respectively. Then

$$\langle T'(x), y \rangle = \langle \lim_{n \rightarrow \infty} T(x_n), \lim_{m \rightarrow \infty} y_m \rangle$$

$$\begin{aligned}
&= \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \langle T(x_n), y_m \rangle \\
&= \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \langle x_n, T^*(y_m) \rangle \\
&= \langle \lim_{n \rightarrow \infty} x_n, \lim_{m \rightarrow \infty} T^*(y_m) \rangle \\
&= \langle x, (T^*)'(y) \rangle.
\end{aligned}$$

(c) Let $x \in \mathcal{H}$, and let $\{w_n\} \subset W$ be a sequence converging to x . Then

$$T'((T^{-1})'(x)) = T'(\lim_{n \rightarrow \infty} T^{-1}(w_n)) = \lim_{n \rightarrow \infty} T'(T^{-1}(w_n)).$$

Since $T^{-1} \in B(W)$ and $w_n \in W$, we have that $T^{-1}(w_n) \in W$ and so $T'(T^{-1}(w_n)) = T(T^{-1}(w_n)) = w_n$. Hence

$$T'((T^{-1})'(x)) = \lim_{n \rightarrow \infty} w_n = x.$$

Conversely, a similar argument shows $(T^{-1})'(T'(x)) = x$, and so $(T^{-1})' = (T')^{-1}$.

□

Proposition 2.2.5. If $T \in BA(W)$ is normal, then its extension $T' \in B(\mathcal{H})$ is normal.

Proof. We need to show that for any $x \in \mathcal{H}$, we have an equality $T'((T')^*(x)) = (T')^*(T'(x))$. By Proposition 2.2.4, this is equivalent to showing $T'((T^*)'(x)) = (T^*)'(T'(x))$. Let $\{w_n\} \subset W$ be a sequence converging to x . Then

$$\begin{aligned}
T'((T^*)'(x)) &= T'(\lim_{n \rightarrow \infty} T^*(w_n)) = \lim_{n \rightarrow \infty} T'T^*(w_n) \\
&= \lim_{n \rightarrow \infty} TT^*(w_n) = \lim_{n \rightarrow \infty} T^*T(w_n) \\
&= \lim_{n \rightarrow \infty} (T^*)'(T'(w_n)) = (T^*)'(\lim_{n \rightarrow \infty} T'(w_n))
\end{aligned}$$

$$=(T^*)'(T'(x)).$$

□

Chapter 3

Spectral Theory

Section 3.1

Introduction

We introduce spectral theory here in the more general context of (not necessarily complete) normed vector spaces X . We write I for the identity operator.

Let $U \subset X$, and let $T : U \rightarrow X$ be a linear operator.

Definition 3.1.1. $\lambda \in \mathbb{C}$ is called *regular* if the following conditions hold:

R1. $(T - \lambda I)^{-1}$ exists.

R2. $(T - \lambda I)^{-1}$ is bounded.

R3. $(T - \lambda I)^{-1}$ has a dense (in X) domain, i.e. $\text{Im}(T - \lambda I)$ is dense in X .

Definition 3.1.2. The set of regular values is called the *resolvent set* of T , and is denoted $\rho(T)$.

Definition 3.1.3. The function R_T mapping $\lambda \mapsto (T - \lambda I)^{-1}$ is called the *resolvent function*.

Definition 3.1.4. The *spectrum* of T is defined as $\sigma(T) := \mathbb{C} \setminus \rho(T)$.

Remark 3.1.5. What we have defined above is the “operator spectrum” as presented in, for example, Chapter 7.2 in [4]. It is apparently defined specifically with regards to a linear operator, and makes no reference to the algebra to which that operator belongs. On the other hand, for a (complex) unital algebra A we can define the “algebraic spectrum” of an element $a \in A$:

$$\sigma_A(a) := \{\lambda \in \mathbb{C} : (x - \lambda 1_A)^{-1} \notin A\},$$

i.e. the set of $\lambda \in \mathbb{C}$ for which $x - \lambda 1_A$ is not invertible in A .

If X is a Banach space, then the operator spectrum of a bounded linear operator $T : X \rightarrow X$ coincides with the algebraic spectrum of $T \in B(X)$.

First suppose λ is regular. Then $(T - \lambda I)^{-1}$ extends uniquely to a bounded linear operator $X \rightarrow X$ by Proposition 2.2.1. Hence $(T - \lambda I)^{-1} \in B(X)$ and $\lambda \notin \sigma_{B(X)}(T)$ according to the definition above. Conversely, suppose λ is such that $(T - \lambda I)^{-1} \in B(X)$. Then immediately λ is regular.

Also for Banach spaces, the resolvent set is sometimes defined as

$$\rho(T) = \{\lambda \in \mathbb{C} : (T - \lambda 1) \text{ is bijective}\}.$$

This is equivalent to Definition 3.1.2. To see this, first suppose $T - \lambda I$ is bijective. Then by the bounded inverse theorem (e.g. 4.12-2 in [4]), we get that $(T - \lambda I)^{-1}$ is also bounded. Thus λ is regular. Conversely, suppose λ is regular. Then as above, $(T - \lambda I)^{-1}$ extends uniquely to a bounded linear operator $X \rightarrow X$, so in particular $(T - \lambda I)$ is bijective.

Remark 3.1.6. For non-Banach spaces, the algebraic and operator spectra may differ. Consider the Hilbert space $\mathcal{H} = L^2[a, b]$ where $0 < a < b$, and consider the proper dense subset of polynomials $P[a, b]$ (it is dense by the Weierstrass approximation theorem, see e.g.

[7]).

Consider the self adjoint multiplication operator $T = M_x \in BA(W) \subset B(\mathcal{H})$. Now $T^{-1} = M_{1/x} \in C[a, b] \subset \mathcal{H}$, however $T^{-1} \notin BA(W)$ since, for example, $\frac{1}{x} \cdot 1 = \frac{1}{x}$ is not a polynomial, hence T^{-1} does not preserve W . This shows that $0 \in \sigma_{BA(W)}(T)$. But we claim $0 \notin \sigma(T)$ with respect to W . It suffices to show 0 is regular (Definition 3.1.1). We have already shown T^{-1} exists, and it is bounded because it is a continuous function with compact domain. Its range is dense by the Müntz-Szász theorem (see again [7]). Thus 0 is regular, hence $0 \notin \sigma(T)$.

In fact, for analagous reasons as the case $\lambda = 0$, we can see that $\sigma_{BA(W)}(T) = \mathbb{C}$. Also, λ fails to be a regular value of $T \in BA(W)$ if and only if $T - \lambda I$ is not invertible, which occurs only when $\lambda \in [a, b]$. The same is true regarding $T \in B(\mathcal{H})$, and so $\sigma(T) = [a, b]$ on both $BA(W)$ and $B(\mathcal{H})$.

The spectrum of a linear operator T can be decomposed as follows:

property of λ			λ belongs to:
R1	R2	R3	
✗	✗	✗	$\sigma_p(T)$
✓	✗	✓	$\sigma_c(T)$
✓	✓	✗	$\sigma_r(T)$
✓	✗	✗	

Remark 3.1.7. R2 and R3 can only be satisfied if R1 is satisfied.

Definition 3.1.8. $\sigma(T)$ can be decomposed into the following disjoint sets:

- the *point spectrum*, denoted $\sigma_p(T)$.

- the *continuous spectrum*, denoted $\sigma_c(T)$.
- the *residual spectrum*, denoted $\sigma_r(T)$.

Definition 3.1.9. Let X be a Banach space, and $T \in B(X)$. A scalar λ is called an *approximate eigenvalue* of T if $T - \lambda I$ is not bounded below, i.e. there exists a sequence of unit vectors (x_n) such that $Tx_n - \lambda x_n \rightarrow 0$. Such a sequence (x_n) is called a *Weyl sequence*. The set of approximate eigenvalues of T is denoted $\sigma_{\text{ap}}(T)$.

Proposition 3.1.10. Approximate eigenvalue lie in the spectrum.

Proof. Let λ be an approximate eigenvalue, and (x_n) a Weyl sequence. If $T - \lambda I$ is not injective, then $\lambda \in \sigma_p(T) \subset \sigma(T)$. So it suffices to consider the case when $T - \lambda I$ is injective. In that case, suppose $(T - \lambda I)^{-1}$ is bounded (hence continuous). But then $\lim_{n \rightarrow \infty} \|x_n\| = 1$ contradicts the computation

$$\begin{aligned} \lim_{n \rightarrow \infty} x_n &= \lim_{n \rightarrow \infty} (T - \lambda I)^{-1} (T - \lambda I)x_n \\ &= (T - \lambda I)^{-1} \lim_{n \rightarrow \infty} (T - \lambda I)x_n \\ &= (T - \lambda I)^{-1}(0) = 0. \end{aligned}$$

In other words, if λ satisfies [R1](#) then it fails [R2](#). Thus $\lambda \in \sigma(T)$. □

Corollary 3.1.11. $\sigma_p(T) \cup \sigma_c(T) \subset \sigma_{\text{ap}}(T)$.

Remark 3.1.12. Every eigenvalue is an approximate eigenvalue by considering the Weyl sequence to be constant, with each term equal to the (normalization of) a chosen eigenvector.

Remark 3.1.13. It is still possible for approximate eigenvalues to be in the residual spectrum. Consider ℓ^2 , and the operator

$$T(x_1, x_2, \dots) = (0, x_1, \frac{x_2}{2}, \frac{x_3}{3}, \dots).$$

We claim $0 \in \sigma_r(T) \cap \sigma_{ap}(T)$.

First let us show that it is even in the spectrum. That is, for $\lambda = 0$, $T - \lambda I = T$ is not invertible. We can explicitly determine that

$$T^{-1}(x_1, x_2, \dots) = (x_2, 2x_3, 3x_4, \dots).$$

This is not bounded, as $\|T^{-1}(e_n)\| = n - 1 \rightarrow \infty$.

Next we will show $0 \in \sigma_r(T)$. We need to show that $0 \notin \sigma_p(T)$ and $0 \notin \sigma_c(T)$. By inspecting its definition we can see that T is injective, so $0 \notin \sigma_p(T)$. Also, $\text{Im}(T) \subset \ell^2$ is not dense. To see this, consider the element $e_1 \notin \text{Im}(T)$. For any $x \in \text{Im}(T)$, we have that $|x - e_1| \geq 1$, and so no sequence in $\text{Im}(T)$ converges to e_1 . Thus $0 \notin \sigma_c(T)$.

Finally, we will show $0 \in \sigma_{ap}(T)$. We calculate that

$$\|T(e_n) - 0 \cdot e_n\| = \|T(e_n)\| = \frac{1}{n} \rightarrow 0,$$

and so $\{e_n\}$ is a Weyl sequence for $\lambda = 0$.

Section 3.2

On Dense Subspaces

Let \mathcal{H} be a separable Hilbert space, and $W \subset \mathcal{H}$ a proper dense subspace. Let $T \in BA(W)$, and let $\tilde{T} \in B(\mathcal{H})$ be its unique extension.

The following are some preliminary observations, some of which will be strengthened in the following sections.

Proposition 3.2.1.

- (a) $\sigma(T) \supset \sigma(\tilde{T})$.

(b) $\sigma_p(T) \subset \sigma_p(\tilde{T})$.

(c) $\sigma_{ap}(T) \subset \sigma_{ap}(\tilde{T})$.

Proof.

(a) We show the contrapositive. Suppose λ is a regular value of T . Then $(T - \lambda I)^{-1}$ exists, is bounded, and its domain is a dense subset of W , hence a dense subset of \mathcal{H} . Thus by Proposition 2.2.4 its unique extension to H is also bounded, and an inverse for $\tilde{T} - \lambda I$. Thus λ is a regular value of \tilde{T} .

(b) If $\lambda \in \sigma_p(T)$ then $T - \lambda I$ is not injective, i.e. there exists $w \in W$ such that $T(w) = \lambda w$. Since \tilde{T} is an extension of T , it must also be the case that $\tilde{T}(w) = \lambda w$, i.e. $\tilde{T} - \lambda I$ is not injective. Hence $\lambda \in \sigma_p(\tilde{T})$.

(c) A Weyl sequence in W for T will also be a Weyl sequence for \tilde{T} .

□

Remark 3.2.2. An operator $T \in BA(W)$ may be injective but have non-injective extension $T \in B(\mathcal{H})$. Consider the Hilbert space ℓ^2 , and let $\{e_i\}$ denote the standard orthonormal basis. Consider the element

$$x_1 = \left(1, \frac{1}{2}, \frac{1}{3}, \dots\right) \in \ell^2.$$

Let $W = \text{span}\{x_1 \cup \{e_i\}_{i \geq 2}\}$. We claim $W \subset \ell^2$ is dense. It suffices to show there is a sequence in W converging (in ℓ^2) to e_1 . Note that

$$\left(x_1 - \sum_{k=2}^i \frac{1}{k} e_k\right)_i = \left(x_1, x_1 - \frac{1}{2}e_2, x_1 - \frac{1}{2}e_2 - \frac{1}{3}e_3, \dots\right)$$

is such a sequence. Now consider the projection operator

$$P(a_1, a_2, a_3, \dots) = (0, a_2, a_3, \dots).$$

Then $\ker(P) = \text{span}\{e_1\}$, which does not lie inside W .

In the following subsections we turn the the question of *spectral permanence*. A deep result of this kind is the following:

Theorem 3.2.3. Let $A \subset B$ be unital C^* -algebras with the same unit, and let $x \in A$. Then $\sigma_A(x) = \sigma_B(x)$.

Proof. See Corollary 3.10 in [9]. □

Remark 3.2.4. Because of this, even if $C^*(W) \neq B(\mathcal{H})$, we at least have “algebraic” spectral permanence. One should be cautious, however, to note that the algebras in the above theorem have the same unit. A situation where this may not be the case is if we consider the unitization of a C^* -subalgebra. For example, the compact operators $K(\mathcal{H})$ are a C^* -subalgebra of $B(\mathcal{H})$ (see, for example, Example 3.3.10 in [9]), the subalgebra contains a unit if and only if \mathcal{H} is finite-dimensional. We may however formally attach a unit in a minimal way while retaining the C^* -structure: this process is called unitization. It is not a priori clear though, that this unit is the same unit in $B(\mathcal{H})$. This turns out to be the case, however, for example because the multiplier algebra of $K(\mathcal{H})$ is $B(\mathcal{H})$ (e.g. Example 13.2.4.2 in [3]).

Section 3.3

Permanence of Approximate Eigenvalues

Proposition 3.3.1. Consider an operator $T \in B(\mathcal{H})$. For any $\lambda \in \sigma_{ap}(T)$, there exists a Weyl sequence for λ completely contained in W .

Proof. By assumption, there exists a Weyl sequence for λ in \mathcal{H} , call it $\{x_n\}$. Since $W \subset \mathcal{H}$ is dense, we can construct a sequence $\{w'_n\}$ such that $\|x_n - w'_n\| < \frac{1}{n}$. Since $\|x_n - w'_n\| \rightarrow 0$, and $\|1 - \|w'_n\|\| \leq \|x_n - w'_n\|$ (by the reverse triangle inequality), it follows that $\|w'_n\| \rightarrow 1$.

We claim the normalization of $\{w'_n\}$, which we will write as $\{w_n\}$, is a Weyl sequence for λ . To see this, let $k > 0$ be such that $\|(T - \lambda I)(x)\| \leq k\|x\|$ for all x (such a k exists since $T - \lambda I$ is bounded). Then:

$$\begin{aligned} \|(T - \lambda I)(w_n)\| &= \frac{1}{\|w'_n\|} \|(T - \lambda I)(w'_n)\| \\ &\leq \frac{1}{\|w'_n\|} (\|(T - \lambda I)(w'_n - a_n)\| + \|(T - \lambda I)(a_n)\|) \\ &\leq \frac{1}{\|w'_n\|} (k\|w'_n - a_n\| + \|(T - \lambda I)(a_n)\|) \\ &< \frac{1}{\|w'_n\|} \left(\frac{k}{n} + \|(T - \lambda I)(a_n)\| \right). \end{aligned}$$

Taking limits, we see that $\|(T - \lambda I)(w_n)\| \rightarrow 0$ as desired. \square

Corollary 3.3.2. Consider an operator $T \in BA(W)$ and its unique extension $\tilde{T} \in B(\mathcal{H})$.

Then $\sigma_{ap}(T) = \sigma_{ap}(\tilde{T})$

Proof. Proposition 3.3.1 implies that $\sigma_{ap}(\tilde{T}) \subset \sigma_{ap}(T)$. The reverse inclusion is discussed in Proposition 3.2.1. \square

Section 3.4

Spectral Permanence for Normal Operators

Proposition 3.4.1. If $T \in B(\mathcal{H})$ is normal, then $\sigma_r(T) = \emptyset$.

Proof. It suffices to show that if $\lambda \in \sigma(T) \setminus \sigma_p(T)$, then $\lambda \in \sigma_c(T)$. So let $\lambda \in \sigma(T) \setminus \sigma_p(T)$.

By Remark 3.1.5, $T - \lambda I$ is injective (yet fails to be surjective). It suffices to show that the $E := \text{Im}(T - \lambda I)$ is dense. Suppose otherwise.

We first observe that there must exist a nonzero $z \in E^\perp$. To see this, note that $\overline{E} \neq \mathcal{H}$ by assumption that E is not dense. We then recall the fact from Hilbert spaces that any

$x \in \mathcal{H}$ can be expressed as $x = y + z$ for $y \in \overline{E}$ and $z \in \overline{E}^\perp$. Thus \overline{E}^\perp is nonempty, and so $E^\perp \supset \overline{E}^\perp$ is also nonempty.

Next, we claim that

$$E^\perp = \text{Ker}(T^* - \overline{\lambda}I) = \text{Ker}(T - \lambda I).$$

This is because of the following facts:

- $(\text{Im}(T))^\perp = \text{Ker}(T^*)$ (Proposition 2.1.4)
- $\text{Ker}(T) = \text{Ker}(T^*)$ (Proposition 2.1.5)

All together, we have asserted the existence of a nonzero $z \in \text{Ker}(T - \lambda I)$, contradicting the fact that T is injective. □

Corollary 3.4.2. For a normal operator $T \in B(\mathcal{H})$, $\sigma(T) = \sigma_{ap}(T)$.

Corollary 3.4.3. Consider an operator $T \in BA(W)$ and its unique extension $\tilde{T} \in B(\mathcal{H})$. If \tilde{T} is normal, then $\sigma_{ap}(T) = \sigma(\tilde{T})$.

Proof. By Corollary 3.4.2, we have that $\sigma(\tilde{T}) = \sigma_{ap}(\tilde{T})$. Then Corollary 3.3.2 implies $\sigma(\tilde{T}) = \sigma_{ap}(T)$. □

Chapter 4

Approximation Results

Section 4.1

Approximation of Compact Operators

Again let \mathcal{H} be a separable Hilbert space, and $W \subset \mathcal{H}$ a proper dense subset. Whenever mentioned, a countable orthonormal basis for \mathcal{H} will be assumed to lie in W (according to Proposition 2.1.3, we lose no generality).

The main goal of this section is to show that every compact operator $T \in \mathcal{H}$ can be approximated by operators in $BA(W)$, i.e. T can be expressed as a limit of operators in $BA(W)$ with respect to the operator norm. Yet another way of viewing this is that $K(\mathcal{H})$ is an ideal of $C^*(W)$.

Definition 4.1.1. An operator T called *compact* if, for every bounded sequence $\{x_n\} \subset \mathcal{H}$, the sequence $\{T(x_n)\}$ contains a convergent subsequence.

Proposition 4.1.2. The following are equivalent:

- (a) T is compact (in the sense of Definition 4.1.1).
- (b) $T(B(0,1))$ is relatively compact, where $B(0,1)$ is the unit ball. In other words,

$\overline{T(B(0,1))}$ is compact.

Proof. See Theorem 8.1-3 in [4]. □

We will write $K(\mathcal{H})$ for the class of compact operators on \mathcal{H} .

Proposition 4.1.3. $K(\mathcal{H})$ is the unique proper closed 2-sided ideal of $B(\mathcal{H})$. It is also a $*$ -ideal.

Proof. See Corollary 5.11 in [2]. □

4.1.1. Finite Rank Operators

Definition 4.1.4. An operator T is called *finite rank* if it is bounded and its image is a finite-dimensional vector space.

We will write $F(\mathcal{H})$ for the set of finite rank operators on \mathcal{H} . We say $T \in F(W)$ if T is finite rank and $T \in B(W)$. Similarly, we write $FA(W)$ for the bounded, adjointable, finite rank operators on W .

Lemma 4.1.5. $F(\mathcal{H}) \subset K(\mathcal{H})$.

Proof. Let $T \in F(\mathcal{H})$, and let $\{x_n\} \subset \mathcal{H}$ be a bounded sequence. Then $\{T(x_n)\}$ is also bounded, and hence must contain a convergent subsequence by the Bolzano-Weierstrass theorem. □

For $x, y \in \mathcal{H}$, we define the operator $\theta_{x,y}$ as follows:

$$\theta_{x,y}(-) = \langle -, x \rangle y.$$

Proposition 4.1.6. If $T \in F(\mathcal{H})$, write $\{e_i\}_1^n$ for an orthonormal basis of $\text{Range}(T)$. Then there exist $\{z_i\}_1^n \in \mathcal{H}$ such that

$$T(x) = \sum_{i=1}^n \langle x, z_i \rangle e_i = \sum_{i=1}^n \theta_{z_i, e_i}(x).$$

Proof. For $\{e_i\}$ as in the statement of the claim, we can write

$$T(x) = \sum_{i=1}^n c_i(x)e_i,$$

where each c_i is a function $\mathcal{H} \rightarrow \mathbb{C}$. First we check that the c_i are linear:

$$\sum_{i=1}^n c_i(x+y)e_i = T(x+y) = T(x) + T(y) = \sum_{i=1}^n (c_i(x) + c_i(y))e_i,$$

which shows that $c_i(x+y) = c_i(x) + c_i(y)$ for all i . Additionally, the c_i are bounded:

$$|c_i(x)| = \|c_i(x)e_i\| \leq \left\| \sum_i c_i(x)e_i \right\| = \|T(x)\| \leq \|T\| \cdot \|x\|.$$

Since the c_i are thus all bounded linear functionals, by the Riesz-Fréchet theorem there exists $z_i \in \mathcal{H}$ such that $c_i(x) = \langle x, z_i \rangle$. In other words,

$$T(x) = \sum_{i=1}^n \langle x, z_i \rangle e_i = \sum_{i=1}^n \theta_{z_i, e_i},$$

as desired. □

Corollary 4.1.7. $F(\mathcal{H}) \subset \text{span}\{\theta_{x,y} : x, y \in \mathcal{H}\}$.

Proposition 4.1.8. $(\theta_{x,y})^* = \theta_{y,x}$

Proof. For $a, b \in \mathcal{H}$,

$$\langle \theta_{x,y}(a), b \rangle = \langle \langle a, x \rangle y, b \rangle = \langle a, x \rangle \cdot \langle y, b \rangle = \langle y, b \rangle \cdot \langle a, x \rangle = \langle a, \langle b, y \rangle x \rangle = \langle a, \theta_{y,x}(b) \rangle.$$

□

Corollary 4.1.9. If $T \in F(\mathcal{H})$, then $\text{Rank}(T) = \text{Rank}(T^*)$.

Proof. By the Propositions 4.1.6 and 4.1.8, we can write

$$T(x) = \sum_{i=1}^n \langle x, z_i \rangle e_i, \quad T^*x = \sum_{i=1}^n \langle x, e_i \rangle z_i,$$

where $\{e_i\}_1^n$ is an orthonormal basis for $\text{Range}(T)$. It suffices to show that the z_i are linearly independent. Suppose otherwise. Without loss of generality, assume $z_1 = \sum_{i=2}^n d_i z_i$. Then $\{e_1 + e_i\}_2^n$ would span the range of T , since

$$\begin{aligned} T(x) &= \sum_{i=1}^n \langle x, z_i \rangle e_i \\ &= \langle x, \sum_{i=2}^n d_i z_i \rangle e_1 + \sum_{i=2}^n \langle x, z_i \rangle e_i \\ &= \sum_{i=2}^n \langle x, (d_i + 1)z_i \rangle (e_1 + e_i). \end{aligned}$$

But the cardinality of the set $\{e_1 + e_i\}_2^n$ contradicts the assumption that $\{e_i\}_1^n$ is a basis for the $\text{Range}(T)$. □

4.1.2. Projection Operators

For $n > 0$, the projection operator $P_n \in B(\mathcal{H})$ with respect to the countable orthonormal basis $\{e_i\}$ maps an element onto its first n components:

$$P_n(x) = P_n \left(\sum_{i=1}^{\infty} \langle x, e_i \rangle e_i \right) = \sum_{i=1}^n \langle x, e_i \rangle e_i.$$

Proposition 4.1.10 (basic facts about P_n).

- (a) $\|P_n\| = 1$.
- (b) $P_n = P_n^*$.
- (c) $P_n \in FA(W)$.

Proof.

(a) Notice that $\|P_n(x)\| \leq \|x\|$ and $\|P_n(e_n)\| = \|e_n\| = 1$, and so $\|P_n\| = 1$.

(b) We calculate:

$$\begin{aligned}\langle P_n x, y \rangle &= \left\langle \sum_{i=1}^n \langle x, e_i \rangle e_i, \sum_{j=1}^{\infty} \langle y, e_j \rangle e_j \right\rangle = \sum_{i=1}^n \langle \langle x, e_i \rangle e_i, \langle y, e_i \rangle e_i \rangle, \\ \langle x, P_n y \rangle &= \left\langle \sum_{j=1}^{\infty} \langle x, e_j \rangle e_j, \sum_{i=1}^n \langle y, e_i \rangle e_i \right\rangle = \sum_{j=1}^n \langle \langle x, e_j \rangle e_j, \langle y, e_j \rangle e_j \rangle.\end{aligned}$$

(c) $\text{Im}(P_n)$ is contained in the span of the $\{e_i\} \subset W$, as is its adjoint by the previous fact. □

Proposition 4.1.11. For any $x \in \mathcal{H}$, we have that $P_n(x) \rightarrow x$.

Proof. We compute

$$\begin{aligned}\|P_n(x) - x\|^2 &= \left\| \sum_{i=1}^{\infty} \langle P_n(x) - x, e_i \rangle e_i \right\|^2 \\ &= \left\| \sum_{i=1}^{\infty} \left\langle \sum_{j=1}^n \langle x, e_j \rangle e_j - \sum_{k=1}^{\infty} \langle x, e_k \rangle e_k, e_i \right\rangle e_i \right\|^2 \\ &= \left\| \sum_{i=1}^{\infty} \left\langle - \sum_{k=n+1}^{\infty} \langle x, e_k \rangle e_k, e_i \right\rangle e_i \right\|^2 \\ &= \left\| - \sum_{i=n+1}^{\infty} \langle x, e_i \rangle e_i \right\|^2 \\ &= \sum_{i=n+1}^{\infty} \|\langle x, e_i \rangle\|^2.\end{aligned}\tag{4.1.12}$$

Pulling the sum out of the norm is justified since all summands are orthogonal. The right hand side tends to 0 as $n \rightarrow \infty$ since $\|x\|$ is finite. □

Lemma 4.1.13. On a compact subset $K \subset \mathcal{H}$, we have that $P_n \rightarrow 1$ uniformly (with respect to the operator norm).

Proof. Since K is compact, we may cover it with finitely many $\sqrt{\epsilon/2}$ -balls. This means that there is a finite set $\{x_1, \dots, x_N\} \subset K$ such that for any $x \in K$ there exists $1 \leq k \leq N$ such that $\|x - x_k\| < \sqrt{\epsilon/2}$.

If we fix $x \in K$ and choose an x_k as above, then, recalling Equation 4.1.12,

$$\begin{aligned}
\|P_n(x) - x\|^2 &= \sum_{i=n+1}^{\infty} \|\langle x, e_i \rangle\|^2 = \sum_{i=n+1}^{\infty} \|\langle x - x_k + x_k, e_i \rangle\|^2 \\
&= \sum_{i=n+1}^{\infty} \|\langle x - x_k, e_i \rangle + \langle x_k, e_i \rangle\|^2 \\
&\leq \sum_{i=n+1}^{\infty} (\|\langle x - x_k, e_i \rangle\| + \|\langle x_k, e_i \rangle\|)^2 \\
&\leq \sum_{i=n+1}^{\infty} \|\langle x - x_k, e_i \rangle\|^2 + \sum_{i=n+1}^{\infty} \|\langle x_k, e_i \rangle\|^2 \\
&\leq \|x - x_k\|^2 + \sum_{i=n+1}^{\infty} \|\langle x_k, e_i \rangle\|^2 \\
&\leq \frac{\epsilon}{2} + \sum_{i=n+1}^{\infty} \|\langle x_k, e_i \rangle\|^2.
\end{aligned}$$

Since there are only finitely many possible x_k , we get that, for any $x \in K$,

$$\|P_n(x) - x\|^2 \leq \frac{\epsilon}{2} + \max_k \sum_{i=n+1}^{\infty} \|\langle x_k, e_i \rangle\|^2.$$

Taking $n \rightarrow \infty$, we get the desired convergence. It does not depend on x , so the convergence is uniform. \square

Corollary 4.1.14. If $T \in K(\mathcal{H})$, then $P_n T \rightarrow T$ uniformly (with respect to the operator norm).

Proof. Since T is compact, $\overline{T(B(0, 1))}$ is compact (by Proposition 4.1.2), where $B(0, 1)$ is

the unit ball. Then we can apply Lemma 4.1.13:

$$\begin{aligned}
\lim_{n \rightarrow \infty} \|P_n T - T\| &= \lim_{n \rightarrow \infty} \sup_{\|x\| \leq 1} \|P_n T(x) - T(x)\| \\
&= \sup_{\|x\| \leq 1} \lim_{n \rightarrow \infty} \|P_n(T(x)) - T(x)\| \\
&= 0,
\end{aligned}$$

where we can interchange the supremum and limit since we are working on the compact subspace $\overline{T(B(0,1))}$, and hence the convergence inside the norm is uniform. Thus, the convergence $P_n T \rightarrow T$ is uniform. \square

Corollary 4.1.15. If $T \in K(\mathcal{H})$, then $TP_n \rightarrow T$ uniformly (with respect to the operator norm).

Proof. In the compact subspace $\overline{B(0,1)}$, we have that $P_n(x) - x \rightarrow 0$ uniformly (by Lemma 4.1.13). Hence $T(P_n(x) - x)$ converges uniformly, and a similar method as above of switching the supremum and limit yields

$$\lim_{n \rightarrow \infty} \|TP_n - T\| = \lim_{n \rightarrow \infty} \sup_{\|x\| \leq 1} \|T(P_n(x) - x)\| = \sup_{\|x\| \leq 1} \lim_{n \rightarrow \infty} \|T(P_n(x) - x)\| = 0.$$

\square

Corollary 4.1.16. If $T \in K(\mathcal{H})$, then $P_n T P_n \rightarrow T$ uniformly (with respect to the operator norm).

Proof. Observe that

$$\begin{aligned}
\|P_n T P_n - T\| &= \|P_n T P_n - P_n T + P_n T - T\| \\
&\leq \|P_n T P_n - P_n T\| + \|P_n T - T\| \\
&= \|P_n\| \cdot \|T P_n - T\| + \|P_n T - T\|.
\end{aligned}$$

By Corollaries 4.1.14 and 4.1.15, we get that $\|P_n T P_n - T\| \rightarrow 0$, so the result follows. (Recall from Proposition 4.1.10 that $\|P_n\| = 1$ for any n .) \square

Corollary 4.1.17. $(P_n T P_n)^* = (P_n T^* P_n)$.

Proof. $(P_n T P_n)^* = P_n^* T^* P_n^*$, and the result follows from the fact $P_n^* = P_n$ (Proposition 4.1.10). \square

Corollary 4.1.18. $P_n T P_n \in FA(W)$.

Proof. This follows from the fact that $P_n \in FA(W)$ (by Proposition 4.1.10) and Corollary 4.1.17. \square

4.1.3. Approximation Result

Corollary 4.1.19. Every $T \in K(\mathcal{H})$ is a limit of finite rank operators on $T_n \in FA(W)$.

Proof. $\{P_n T P_n\} \subset FA(W)$ (by Corollary 4.1.18) converges to T (by Corollary 4.1.16). \square

Proposition 4.1.20. $K(\mathcal{H})$ is an ideal of $C^*(W)$.

Proof. $K(\mathcal{H}) \subset C^*(W)$ by the previous result, and the ideal structure follows immediately from the fact that $K(\mathcal{H})$ is an ideal of $B(\mathcal{H})$ (Proposition 4.1.3). \square

Section 4.2

Operators on the Span of an ON basis

Now consider the subspace $W_0 = \text{span}\{e_n\}$. Then $W_0 \subset W \subset \mathcal{H}$ and $W_0 \subset \mathcal{H}$ is dense.

4.2.1. Compatibility $BA(W_0)$ and $BA(W)$

Since $BA(W) \subset BA(\mathcal{H}) = B(\mathcal{H})$, it is natural to ask whether $BA(-)$ is “functorial” with respect to inclusions. For example: is it true that $BA(W_0) \subset BA(W)$, or $BA(W) \subset BA(W_0)$? The answer in both cases, interestingly, is not necessarily.

Example 4.2.1 ($BA(W) \not\subset BA(W_0)$). Let $\mathcal{H} = L^2[0, 1]$. Let $W = C[0, 1]$ and $W_0 = \text{span}\{1, x, x^2, \dots\}$. It is also well known that $W \subset \mathcal{H}$ is dense (e.g. Consequence (i) of Lemma 4.9 in [6]). $W_0 \subset W$ is dense by the Weierstrass approximation theorem, hence W_0 is a span of a (countable) orthonormal basis, e.g. the Hermite polynomials. Thus we satisfy the conditions we want.

Consider the multiplication operator M_{e^x} . It preserves W since the product of two continuous functions is continuous, and is also self-adjoint, so $M_{e^x} \in BA(W)$. However, $1 \cdot e^x = e^x$ is not in the span of the polynomials, and so $M_{e^x} \notin BA(W_0)$.

Example 4.2.2 ($BA(W_0) \not\subset BA(W)$). Let \mathcal{H} and W_0 be as in the previous example, but now let

$$W = \text{span}(W_0 \cup \{\text{simple functions in } \mathcal{H}\}).$$

W_0 is already dense in \mathcal{H} by our previous remarks, so W is as well and we satisfy the condition we want.

But consider the self-adjoint multiplication operator M_x . Then $M_x \in BA(W_0)$. But consider the simple function $f = \mathbf{1}_{[0, 1/3]} + \mathbf{1}_{[2/3, 1]}$. Then $xf(x)$ is neither simple nor continuous (and since it is not continuous, it is not spanned by the polynomials). Hence $M_x \notin BA(W)$.

4.2.2. Density of $BA(W_0) \subset B(\mathcal{H})$

The tractable linear structure of W_0 allows us to prove the following result, due to [8]. The idea is to show that, given any $T \in B(\mathcal{H})$, we can find a $B \in BA(W_0)$ that is arbitrarily close to T (with respect to the operator norm).

Fix a nonzero $A \in B(\mathcal{H})$. Let $Q_n := I - P_n$, i.e.

$$Q_n(x) = \sum_{i=n+1}^{\infty} \langle x, e_i \rangle e_i.$$

Since $A(e_n)$ and $A^*(e_n)$ have finite norm, for any $\epsilon > 0$ and $n \geq 1$ we can find $k_n > n$ such

that

$$\|Q_{k_n}A(e_n)\| \leq \frac{\epsilon}{3 \cdot 2^{n/2}} \quad \text{and} \quad \|Q_{k_n}A^*(e_n)\| \leq \frac{\epsilon}{3 \cdot 2^{n/2}}.$$

Now define the operator $R \in B(\mathcal{H})$ as follows:

$$R(x) = \sum_{n=1}^{\infty} \left(\langle x, e_n \rangle Q_{k_n}A(e_n) + \langle x, Q_{k_n}A^*(e_n) \rangle e_n \right).$$

Proposition 4.2.3. $R \in B(\mathcal{H})$.

Proof. By the Cauchy-Schwartz inequality, we have on the one hand

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \langle x, e_n \rangle Q_{k_n}A(e_n) \right\| &\leq \sum_{n=1}^{\infty} \|\langle x, e_n \rangle Q_{k_n}A(e_n)\| = \sum_{n=1}^{\infty} |\langle x, e_n \rangle| \cdot \|Q_{k_n}A(e_n)\| \\ &\leq \sum_{n=1}^{\infty} \|x\| \cdot \frac{\epsilon}{3 \cdot 2^{n/2}} \leq \|x\| \left(\sum_{n=1}^{\infty} \left(\frac{\epsilon}{3 \cdot 2^{n/2}} \right)^2 \right)^{1/2} \\ &= \frac{\epsilon \|x\|}{3} \cdot \left(\sum_{n=1}^{\infty} 2^{-n} \right)^{1/2} = \frac{\epsilon \|x\|}{3}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \langle x, Q_{k_n}A^*(e_n) \rangle e_n \right\| &= \left(\sum_{n=1}^{\infty} |\langle x, Q_{k_n}A^*(e_n) \rangle| \right)^{1/2} \leq \|x\| \cdot \left(\sum_{n=1}^{\infty} \|Q_{k_n}A^*(e_n)\| \right)^{1/2} \\ &\leq \frac{\epsilon \|x\|}{3} \end{aligned}$$

by an analogous argument. Combining these inequalities shows that R is bounded:

$$\begin{aligned} \|R(x)\| &= \left\| \sum_{n=1}^{\infty} \left(\langle x, e_n \rangle Q_{k_n}A(e_n) + \langle x, Q_{k_n}A^*(e_n) \rangle e_n \right) \right\| \\ &\leq \left\| \sum_{n=1}^{\infty} \langle x, e_n \rangle Q_{k_n}A(e_n) \right\| + \left\| \sum_{n=1}^{\infty} \langle x, Q_{k_n}A^*(e_n) \rangle e_n \right\| \\ &\leq \frac{\epsilon \|x\|}{3} + \frac{\epsilon \|x\|}{3} \end{aligned}$$

$$\langle \epsilon \|x\| \rangle.$$

□

Proposition 4.2.4.

$$R^*(x) = \sum_{n=1}^{\infty} (\langle x, Q_{k_n} A(e_n) \rangle e_n + \langle x, e_n \rangle Q_{k_n} A^*(e_n)).$$

Proof. One the one hand,

$$\begin{aligned} \left\langle \sum_{n=1}^{\infty} \langle x, e_n \rangle Q_{k_n} A(e_n), y \right\rangle &= \sum_{n=1}^{\infty} \langle x, e_n \rangle \cdot \langle Q_{k_n} A(e_n), y \rangle \\ &= \sum_{n=1}^{\infty} \langle x, \langle y, Q_{k_n} A(e_n) \rangle e_n \rangle \end{aligned}$$

and so

$$\left(\sum_{n=1}^{\infty} \langle -, e_n \rangle Q_{k_n} A(e_n) \right)^* = \sum_{n=1}^{\infty} \langle -, Q_{k_n} A(e_n) \rangle e_n.$$

On the other hand,

$$\begin{aligned} \left\langle \sum_{n=1}^{\infty} \langle x, Q_{k_n} A^* e_n \rangle e_n, y \right\rangle &= \sum_{n=1}^{\infty} \langle x, Q_{k_n} A^* e_n \rangle \cdot \langle e_n, y \rangle \\ &= \sum_{n=1}^{\infty} \langle x, \langle y, e_n \rangle Q_{k_n} A^* e_n \rangle \end{aligned}$$

and so

$$\left(\sum_{n=1}^{\infty} \langle -, Q_{k_n} A^* e_n \rangle e_n \right)^* = \sum_{n=1}^{\infty} \langle -, e_n \rangle Q_{k_n} A^*(e_n).$$

Summing the two gives the desired result. □

Theorem 4.2.5. $BA(W_0) \subset B(\mathcal{H})$ is dense.

Proof. Let $B = A - R$. The fact that $B \in B(\mathcal{H})$ follows from Proposition 4.2.3. We claim

that $\|A - B\| \leq \epsilon$ and $B \in BA(W_0)$. With regards to the first claim,

$$\|A - B\| = \|R\| < \epsilon$$

by the previous calculation that R is bounded.

To show $B \in BA(W_0)$, we first show $B(W_0) \subset W_0$. Since W_0 is the span of the $\{e_n\}$, it suffices to show that $B(e_n) \in W_0$ for all n :

$$\begin{aligned} B(e_n) &= A(e_n) - R(e_n) \\ &= A(e_n) - \left(\sum_{i=1}^{\infty} (\langle e_n, e_i \rangle Q_{k_i} A(e_i) + \langle e_n, Q_{k_i} A^*(e_i) \rangle e_i) \right) \\ &= A(e_n) - Q_{k_n} A(e_n) - \sum_{i=1}^{n-1} \langle e_n, Q_{k_i} A^*(e_i) \rangle e_i \\ &= (I - Q_{k_n}) A(e_n) - \sum_{i=1}^{n-1} \langle e_n, Q_{k_i} A^*(e_i) \rangle e_i \\ &= P_{k_n} A(e_n) - \sum_{i=1}^{n-1} \langle e_n, Q_{k_i} A^*(e_i) \rangle e_i \in W_0, \end{aligned}$$

where in the third line the sum cuts off at n because Q_{k_i} projects onto the $\{e_{j > k_i}\}$, hence will vanish when taking an inner product with against an $e_{m \leq k_i}$.

All that remains is to show $B^*(W_0) \subset W_0$. Using Proposition 4.2.4, we carry out an analogous computation:

$$\begin{aligned} B^*(e_n) &= A^*(e_n) - R^*(e_n) \\ &= A^*(e_n) - \left(\sum_{i=1}^{\infty} (\langle e_n, Q_{k_i} A(e_i) \rangle e_i + \langle e_n, e_i \rangle Q_{k_i} A^*(e_i)) \right) \\ &= A^*(e_n) - Q_{k_n} A^*(e_n) - \sum_{i=1}^{n-1} \langle e_n, Q_{k_i} A^*(e_i) \rangle e_i \end{aligned}$$

$$= P_{k_n} A^*(e_n) - \sum_{i=1}^{n-1} \langle e_n, Q_{k_i} A^*(e_i) \rangle e_i \in W_0.$$

Hence $B \in BA(W_0)$. □

Section 4.3

Example: $BA(W) \subset B(\mathcal{H})$ not Dense

We have seen that for proper dense subspaces $W_0 \subset \mathcal{H}$ which are linear spans of an orthonormal basis, it is true that $BA(W_0) \subset BA(\mathcal{H})$ is dense. However, we have also seen that, given an arbitrary proper dense subspace $W \subset \mathcal{H}$ and the linear span of an orthonormal basis contained in it $W_0 \subset W$, operators which preserve W_0 need not preserve W , and vice versa. Thus we might be pessimistic as to whether $BA(W) \subset B(\mathcal{H})$ is dense in general. The purpose of this section is to provide such a counterexample.

The setup is the following: let $\mathcal{H} = \ell^2$ and $W = \ell^1$ ($W \subset \mathcal{H}$ is dense because, for example, the set of eventually 0 sequences is in both of them). The idea is to construct $V \in B(\mathcal{H})$ such that $\|T - V\| > 1/2$ for all $T \in BA(W)$. In the following, $\{e_i\}_1^\infty$ will denote the canonical orthonormal basis for ℓ^2 .

The proof idea is due to [1], but executed and corrected here in full detail, in recognition of the necessity of considering bounded *adjointable* operators on the subspace rather than simply bounded operators.

4.3.1. An Orthogonal Sequence: (ξ_n)

Let

$$\xi_n := 2^{-n/2} \sum_{i=0}^{2^n-1} e_{2^n+i}.$$

Remark 4.3.1. We write the first few ξ_n explicitly:

$$\xi_1 = 2^{-1/2} \cdot (0, 1, 1, 0, \dots)$$

$$\xi_2 = 2^{-1} \cdot (0, 0, 0, 1, 1, 1, 1, 0, \dots)$$

$$\xi_3 = 2^{-3/2} \cdot (0, 0, 0, 0, 0, 0, 0, 1, 1, 1, 1, 1, 1, 1, 0, \dots)$$

Note $2^{-1/2} > 2^{-1} > 2^{-3/2} > \dots$, i.e. $2^{-n/2}$ is a monotonically decreasing sequence.

Proposition 4.3.2. The ξ_n are mutually orthogonal in ℓ^2 , i.e. $\langle \xi_n, \xi_m \rangle = 0$ whenever $n \neq m$.

Proof. Assume, without loss of generality, that $n < m$. Then $2^n + i < 2^m$ for any $0 \leq i \leq 2^n - 1$. (Indeed, even if $m = n + 1$, the closest these values get is $2^n + (2^n - 1) = 2^{n+1} - 1 = 2^m - 1 < 2^m$.) We thus proceed as follows:

$$\begin{aligned} \left\langle 2^{-n/2} \sum_{i=0}^{2^n-1} e_{2^n+i}, 2^{-m/2} \sum_{j=0}^{2^m-1} e_{2^m+j} \right\rangle &= 2^{-n/2} \sum_{i=0}^{2^n-1} \left\langle e_{2^n+i}, 2^{-m/2} \sum_{j=0}^{2^m-1} e_{2^m+j} \right\rangle \\ &= 2^{-n/2} \sum_{i=0}^{2^n-1} \left(\overline{2^{-m/2} \sum_{j=0}^{2^m-1} \langle e_{2^m+j}, e_{2^n+i} \rangle} \right) \\ &= 2^{-n/2} \sum_{i=0}^{2^n-1} \left(\overline{2^{-m/2} \sum_{j=0}^{2^m-1} 0} \right) \\ &= 0. \end{aligned}$$

□

Proposition 4.3.3. For all n ,

$$\|\xi_n\|_p = \begin{cases} \left(2^{\frac{n(2-p)}{2}} \right)^{1/p} & 1 \leq p < \infty \\ 2^{-n/2} & p = \infty \end{cases}.$$

Proof. Note

$$\begin{aligned}
 \xi_n &= 2^{-n/2} \sum_{i=0}^{2^n-1} e_{2^n+i} \\
 &= 2^{-n/2} \cdot (\dots, 0, \underbrace{1, \dots, 1}_{2^n \text{ copies}}, 0, \dots) \\
 &= (\dots, 0, \underbrace{2^{-n/2}, \dots, 2^{-n/2}}_{2^n \text{ copies}}, 0, \dots),
 \end{aligned}$$

and so

$$\|\xi_n\|_p^p = 2^n \cdot (2^{-n/2})^p = 2^{\frac{n(2-p)}{2}}$$

for $1 \leq p < \infty$, and $\|\xi_n\|_\infty = 2^{-n/2}$. □

Corollary 4.3.4.

$$\begin{aligned}
 \|\xi_n\|_1 &= 2^{n/2} \\
 \|\xi_n\|_2 &= 1
 \end{aligned}$$

Corollary 4.3.5.

$$\lim_{n \rightarrow \infty} \|\xi_n\|_p = \begin{cases} \infty & 1 \leq p < 2, \\ 1 & p = 2, \\ 0 & p > 2 \end{cases}.$$

4.3.2. The Operator V

Now define the operator

$$V(x) := V \left(\sum_{n=1}^{\infty} \langle x, e_n \rangle e_n \right) = \sum_{n=1}^{\infty} \langle x, e_n \rangle \xi_n$$

for all $x \in \ell^2$.

Proposition 4.3.6. $V(e_i) = \xi_i$ for all i .

Proof.

$$V(e_i) = \sum_{n=1}^{\infty} \langle e_i, e_n \rangle \xi_n = \langle e_i, e_i \rangle \xi_i = \xi_i.$$

□

Proposition 4.3.7. V is a (partial) isometry. In particular, $V \in B(\ell^2)$.

Proof. For any $x \in \ell^2$,

$$\begin{aligned} \|V(x)\|_2^2 &= \left\| \sum_{n=1}^{\infty} \langle x, e_n \rangle \xi_n \right\|_2^2 \\ &= \sum_{n=1}^{\infty} \|\langle x, e_n \rangle \xi_n\|_2^2, \end{aligned}$$

by the Pythagorean theorem, since the ξ_n are mutually orthonormal,

$$\begin{aligned} &= \sum_{n=1}^{\infty} |\langle x, e_n \rangle|^2 \\ &= \|x\|_2^2 \end{aligned}$$

by Parseval's identity.

□

Proposition 4.3.8. For $T \in B(W)$, if $\|T - V\|_2 < 1/2$, then

$$\sum_{i=0}^{2^n-1} |\langle e_{2^n+i}, T e_n \rangle| \geq \frac{2^{n/2}}{2}.$$

Proof. We first show that

$$\sum_{i=0}^{2^n-1} |\langle e_{2^n+i}, T(e_n) \rangle| \geq 2^{n/2} - \left| \sum_{i=0}^{2^n-1} (\langle e_{2^n+i}, T(e_n) \rangle - 2^{-n/2}) \right|.$$

To see this, first observe that

$$\begin{aligned} \left| \sum_{i=0}^{2^n-1} (\langle e_{2^n+i}, T(e_n) \rangle - 2^{-n/2}) \right| &= \left| \left(\sum_{i=0}^{2^n-1} \langle e_{2^n+i}, T(e_n) \rangle \right) - 2^{n/2} \right| \\ &\geq \left| \left| \sum_{i=0}^{2^n-1} \langle e_{2^n+i}, T(e_n) \rangle \right| - 2^{n/2} \right| \\ &= \left| 2^{n/2} - \left| \sum_{i=0}^{2^n-1} \langle e_{2^n+i}, T(e_n) \rangle \right| \right| \\ &\geq 2^{n/2} - \left| \sum_{i=0}^{2^n-1} \langle e_{2^n+i}, T(e_n) \rangle \right|. \end{aligned}$$

Therefore,

$$\begin{aligned} 2^{n/2} - \left| \sum_{i=0}^{2^n-1} (\langle e_{2^n+i}, T(e_n) \rangle - 2^{-n/2}) \right| &\leq 2^{n/2} - \left(2^{n/2} - \left| \sum_{i=0}^{2^n-1} \langle e_{2^n+i}, T(e_n) \rangle \right| \right) \\ &= \left| \sum_{i=0}^{2^n-1} \langle e_{2^n+i}, T(e_n) \rangle \right| \\ &\leq \sum_{i=0}^{2^n-1} |\langle e_{2^n+i}, T(e_n) \rangle| \end{aligned}$$

as desired.

Now we will show that

$$2^{-n/2} \left| \sum_{i=0}^{2^n-1} (\langle e_{2^n+i}, T(e_n) \rangle - 2^{-n/2}) \right| \leq \|T(e_n) - \xi_n\|_2.$$

We calculate

$$\begin{aligned}
2^{-n/2} \left| \sum_{i=0}^{2^n-1} (\langle e_{2^n+i}, T(e_n) \rangle - 2^{-n/2}) \right| &= \left| \sum_{i=0}^{2^n-1} (2^{-n/2} \langle e_{2^n+i}, T(e_n) \rangle - 2^{-n}) \right| \\
&= |\langle \xi_n, T(e_n) \rangle - 1| \\
&= |\langle \xi_n, T(e_n) \rangle - \langle \xi_n, \xi_n \rangle| \\
&= |\langle \xi_n, T(e_n) - \xi_n \rangle| \\
&\leq \|T(e_n) - \xi_n\|_2,
\end{aligned}$$

where we have used the Cauchy-Schwartz inequality and the fact that $\|\xi_n\|_2 = 1$.

We also claim that

$$\|T(e_n) - \xi_n\|_2 \leq 1/2.$$

This is because $\|T - V\|_2 < 1/2$ by assumption, and since e_n is unit size (in the 2-norm), we get from the properties of the norm that $\|T(e_n) - V(e_n)\|_2 = \|T(e_n) - \xi_n\|_2 < 1/2$.

Pulling it all together, we calculate that

$$\begin{aligned}
\sum_{i=0}^{2^n-1} |\langle e_{2^n+i}, T(e_n) \rangle| &\geq 2^{n/2} - \left| \sum_{i=0}^{2^n-1} (\langle e_{2^n+i}, T(e_n) \rangle - 2^{-n/2}) \right| \\
&= 2^{n/2} \left(1 - 2^{-n/2} \left| \sum_{i=0}^{2^n-1} (\langle e_{2^n+i}, T(e_n) \rangle - 2^{-n/2}) \right| \right) \\
&\geq 2^{n/2} (1 - \|T(e_n) - \xi_n\|_2) \\
&\geq 2^{n/2} \left(1 - \frac{1}{2} \right) \\
&= \frac{2^{n/2}}{2}.
\end{aligned}$$

□

4.3.3. Two Sequences

Proposition 4.3.9. There exist two nonnegative, monotonically increasing sequences of integers $(\alpha)_0^\infty, (\beta)_1^\infty$ such that

$$\sum_{i \notin [\alpha_{r-1}, \alpha_r - 1]} |\langle e_i, T(e_{\beta_r}) \rangle| < 1$$

for all $r \in \mathbb{N}$. We can pick these such that $\alpha_0 = \beta_1 = 1$ and $\beta_i \geq i^2$.

Proof. We will construct the two sequences inductively. Let $\alpha_0 = \beta_1 = 1$. Since $e_{\beta_r} \in \ell^1$, $T(e_{\beta_r}) \in \ell^1$ and

$$\|T(e_{\beta_r})\|_1 = \sum_{i=1}^{\infty} |\langle e_i, T(e_{\beta_r}) \rangle| < \infty.$$

Therefore, there must exist $\alpha_1 > 1$ such that all coordinates of $T(e_{\beta_r})$ outside the range $[\alpha_0, \alpha_1]$ sum to less than 1. In other words, there must be an $\alpha_1 > \alpha_0$ such that

$$\sum_{i \notin [\alpha_0, \alpha_1]} |\langle e_i, T(e_{\beta_r}) \rangle| < 1.$$

For the induction step, suppose we have constructed $(\alpha)_0^r$ and $(\beta)_1^r$. First we will construct β_{r+1} . Note that $e_q \in \ell^1$ for any $q \in \mathbb{N}$, so $T(e_q)$ and $T^*(e_q)$ are both in ℓ^1 . Thus

$$\sum_{i=1}^{\alpha_r - 1} |\langle e_i, T(e_q) \rangle| = \sum_{i=1}^{\alpha_r - 1} |\langle T^*(e_i), e_q \rangle|.$$

The expression $\langle T^*(e_i), e_q \rangle$ picks out the q th coordinate of an element in ℓ^1 . Therefore, we may pick $q = \beta_{r+1} > \beta_r$ such that

$$\max_{i \in [1, \alpha_r - 1]} |\langle T^*(e_i), e_q \rangle| < \frac{1}{2\alpha_r}.$$

This will ensure that

$$\sum_{i=1}^{\alpha_r-1} |\langle e_i, T(e_{\beta_{r+1}}) \rangle| < \frac{1}{2}.$$

Nothing prevents us from picking such that $\beta_{r+1} \geq (r+1)^2$ as well.

Outside of $[1, \alpha_r - 1]$, which is the range $[\alpha_r, \infty)$, the sum is still finite. Thus we can once again find an upper bound $\alpha_{r+1} > \alpha_r$ such that the above sum is $< 1/2$ on $[\alpha_{r+1}, \infty)$. Pulling this all together, we have found $\alpha_{r+1} > \alpha_r$ such that

$$\sum_{i \notin [\alpha_r, \alpha_{r+1}-1]} |\langle e_i, T(e_{\beta_{r+1}}) \rangle| < 1.$$

This completes the induction step. □

4.3.4. Failure to Approximate V

Theorem 4.3.10. Taking $\mathcal{H} = \ell^2$ and $W = \ell^1$ as above, $BA(W) \subset B(\mathcal{H})$ is not dense.

Proof. Suppose there exists $T \in BA(\ell^1)$ such that $\|T - V\|_2 < 1/2$. Consider the element

$$w = \sum_{n=1}^{\infty} \frac{1}{n^2} e_{\beta_n} \in \ell^1.$$

This element is indeed in ℓ^1 as $\sum_n 1/n^2 = \pi/6 < \infty$.

Our first claim is that the series

$$\sum_{n=1}^{\infty} \left(\frac{2^{\beta_n/2}}{2n^2} - \frac{\pi^2}{6} \right)$$

diverges. Indeed,

$$\lim_{n \rightarrow \infty} \left(\frac{2^{\beta_n/2}}{2n^2} - \frac{\pi^2}{6} \right) \rightarrow \infty$$

since in particular $\beta_n \geq n^2$.

Now

$$\begin{aligned}
\|Tw\|_1 &= \sum_i \left| \left\langle e_i, \sum_k \frac{1}{k^2} T(e_{\beta_k}) \right\rangle \right| \\
&= \sum_n \sum_{i=\alpha_{n-1}}^{\alpha_n-1} \left| \left\langle e_i, \sum_k \frac{1}{k^2} T(e_{\beta_k}) \right\rangle \right| \\
&= \sum_n \sum_{i=\alpha_{n-1}}^{\alpha_n-1} \left| \frac{1}{n^2} \langle e_i, T(e_{\beta_n}) \rangle + \sum_{k \neq n} \frac{1}{k^2} \langle e_i, T(e_{\beta_k}) \rangle \right| \\
&\geq \sum_n \sum_{i=\alpha_{n-1}}^{\alpha_n-1} \left(\left| \frac{1}{n^2} \langle e_i, T(e_{\beta_n}) \rangle \right| - \left| \sum_{k \neq n} \frac{1}{k^2} \langle e_i, T(e_{\beta_k}) \rangle \right| \right),
\end{aligned}$$

by an application of the triangle inequality,

$$\begin{aligned}
&\geq \sum_n \sum_{i=\alpha_{n-1}}^{\alpha_n-1} \left(\frac{1}{n^2} |\langle e_i, T(e_{\beta_n}) \rangle| - \sum_{k \neq n} \frac{1}{k^2} |\langle e_i, T(e_{\beta_k}) \rangle| \right) \\
&= \sum_n \left(\sum_{i=\alpha_{n-1}}^{\alpha_n-1} \frac{1}{n^2} |\langle e_i, T(e_{\beta_n}) \rangle| - \sum_{i=\alpha_{n-1}}^{\alpha_n-1} \sum_{k \neq n} \frac{1}{k^2} |\langle e_i, T(e_{\beta_k}) \rangle| \right) \\
&= \sum_n \left(\sum_{i=\alpha_{n-1}}^{\alpha_n-1} \frac{1}{n^2} |\langle e_i, T(e_{\beta_n}) \rangle| - \sum_{k \neq n} \frac{1}{k^2} \sum_{i=\alpha_{n-1}}^{\alpha_n-1} |\langle e_i, T(e_{\beta_k}) \rangle| \right),
\end{aligned}$$

by Tonelli's theorem for series,

$$\geq \sum_n \left(\sum_{i=\alpha_{n-1}}^{\alpha_n-1} \frac{1}{n^2} |\langle e_i, T(e_{\beta_n}) \rangle| - \sum_{k \neq n} \frac{1}{k^2} \right),$$

by Proposition 4.3.9, since $k \neq n$,

$$\begin{aligned}
&= \sum_n \left(\sum_{i=\alpha_{n-1}}^{\alpha_n-1} \frac{1}{n^2} |\langle e_i, T(e_{\beta_n}) \rangle| - \frac{\pi^2}{6} + \frac{1}{n^2} \right) \\
&= \sum_n \left(\frac{1}{n^2} \sum_{i=\alpha_{n-1}}^{\alpha_n-1} |\langle e_i, T(e_{\beta_n}) \rangle| - \frac{\pi^2}{6} + \frac{1}{n^2} \right)
\end{aligned}$$

$$\geq \sum_n \left(\frac{1}{n^2} \left(\sum_i |\langle e_i, T(e_{\beta_n}) \rangle| - 1 \right) - \frac{\pi^2}{6} + \frac{1}{n^2} \right),$$

by Proposition 4.3.9,

$$\begin{aligned} &\geq \sum_n \left(\frac{1}{n^2} \sum_i |\langle e_i, T(e_{\beta_n}) \rangle| - \frac{\pi^2}{6} \right) \\ &\geq \sum_n \left(\frac{1}{n^2} \sum_{i=2^{\beta_n}}^{2^{\beta_n+1}-1} |\langle e_i, T(e_{\beta_n}) \rangle| - \frac{\pi^2}{6} \right) \\ &= \sum_n \left(\frac{1}{n^2} \sum_{i=0}^{2^{\beta_n}-1} |\langle e_{2^{\beta_n}+i}, T(e_{\beta_n}) \rangle| - \frac{\pi^2}{6} \right) \\ &\geq \sum_n \left(\frac{2^{\beta_n/2}}{2n^2} - \frac{\pi^2}{6} \right), \end{aligned}$$

by Proposition 4.3.8,

$$= \infty.$$

This shows that $\|T(w)\|_1$ is bounded below by a divergent series, hence $\|T(w)\|_1$ is also divergent. This slight abuse of notation nonetheless demonstrates that $T(w) \notin \ell^1$. But this is a contradiction since $T \in BA(\ell^1)$. Hence $\|T - V\| > 1/2$ for all $T \in BA(\ell^1)$. \square

Chapter 5

Future Directions

We have shown $BA(\ell^1) \subset B(\ell^2)$ is not dense, and would like to extend these results to ℓ^p for $1 < p < 2$. We conjecture that $BA(\ell^p) \subset B(\mathcal{H})$ is dense. In the case of ℓ^1 , we have the following nontrivial extension:

$$0 \rightarrow K(\ell^2) \rightarrow C^*(\ell^1) \rightarrow \frac{C^*(\ell^1)}{K(\ell^2)} \rightarrow 0.$$

One could apply K -theory to this sequence to see whether the K -theory detects the fact that $C^*(\ell^1) \neq B(\ell^2)$. More generally, one can compare properties of the Calkin algebra $B(\ell^2)/K(\ell^2)$ to the properties of this distinct algebra $C^*(\ell^1)/K(\ell^2)$.

Another direction could be to consider weaker topologies. For the span of an orthonormal basis W_0 , is it still true that $BA(W_0) \subset B(\mathcal{H})$ is dense with respect to the strong/weak operator topologies? Furthermore, it is known that the commutant of a self-adjoint subspace of operators is a von Neumann algebra, so $BA(W)'$ is a von Neumann algebra. One could compute this, and, if it turns out to be nontrivial, apply methods of W^* -algebras to $BA(W)'$. One could also approach the question of density of $BA(W)$ by computing the double commutant $BA(W)''$; the theory shows that if $BA(W) = BA(W)''$ then $BA(W)$ is closed in the strong/weak operator topologies, hence closed in the operator norm topology.

Then if $BA(W) \subsetneq B(\mathcal{H})$, it is certainly not dense.

Finally, one could expand on the spectral theory of $BA(W)$ developed here, or consider the spectral theory of a nontrivial $C^*(W)$. How much of the spectral theory on $B(\mathcal{H})$ can be recovered, and does the spectrum/resolvent “see” certain properties of $BA(W)$?

Bibliography

- [1] Yemon Choi and heller, *Bounded operators leaving dense subspace invariant*, Jun 2017.
- [2] Ronald G. Douglas, *Banach algebra techniques in operator theory*, Springer, 1998.
- [3] Ilijas Farah, *Combinatorial set theory of C^* -algebras*, Springer, 2019.
- [4] Erwin Kreyszig, *Introductory functional analysis with applications*, Wiley, 1989.
- [5] Peter D. Lax, *Functional analysis*, Wiley, 2002.
- [6] Jean-François LeGall, *Measure theory, probability, and stochastic processes*, Springer International Publishing, 2022.
- [7] Andreu Ferré Moragues, *What is... the Müntz–Szász theorem?*
- [8] Ryszard Szwarc, *Algebras of adjointable operators on dense subspaces of Hilbert space*, Jun 2022.
- [9] Ivan F Wilde, *An introduction to C^* -algebras*.
- [10] Neil Young, *An introduction to Hilbert space*, Cambridge University Press, 1988.